

Cyclosporin Reference Range:

Therapeutic range based on draw immediately prior to next dose. Therapeutic Default Reference Range: 50 - 400 ug/L	
Kidney Transplant (months post transplant):	Range (mg/ml)
Pediatric	
0-3 months:	175 - 200
3 - 6 months:	150 - 175
6 - 9 months:	125 - 150
9 - 12 months:	100 - 125
>12 months:	75 - 100
Adult	
0-3 months:	175 - 200
3 - 6 months:	150 - 200
6 - 12 months:	125 - 150
>12 months:	75 - 100
Heart Transplant (months post transplant)	Range (mg/ml)
Pediatric	
0- 12 months:	200 -250
>12 months:	100 - 125
EBV peR>1000 copies/mL:	80 - 100
Adult	
0-3 months:	150 - 250
3 - 6 months:	125 - 225
6 - 12 months:	100 - 200
>12 months:	50 - 150
Heart - Lung Block	200 - 250
Lung Transplant (months post transplant)	Range (mg/ml)
0- 12 months:	175 - 225
>12 months:	125 -175
Liver Transplant (months post transplant)	Range (mg/ml)
0-6 months:	150 - 200
>6 months:	100 - 150
Pancreas Transplant (months post transplant)	Range (mg/ml)
0-6 months:	200 - 250
6 - 12 months:	150 - 200
>12 months:	100 - 150
Bone Marrow Transplant	200 - 400 mg/ml

Critical Values: >400 ug/L

Limitations: The recommended therapeutic ranges described above apply to trough Specimens collected just before the next dose. Blood collected at other times will yield higher results.